

MASS SPECTROMETRIC ANALYSIS OF LONG-CHAIN LIPIDS

Robert C. Murphy^{1*} and Paul H. Axelsen^{2**}

¹Department of Pharmacology, University of Colorado Denver, MSC 8303, 12801 E. 17th Avenue, Aurora, CO 80045

²Departments of Pharmacology, Biochemistry and Biophysics and Medicine, University of Pennsylvania School of Medicine, Room 105 Johnson Pavilion, 3610 Hamilton Walk, Philadelphia, PA 19104-6084

Published online 8 November 2010 in Wiley Online Library (wileyonlinelibrary.com) DOI 10.1002/mas.20284

Electrospray and matrix assisted laser desorption ionization generate abundant molecular ion species from all known lipids that have long chain fatty acyl groups esterified or amidated to many different polar headgroup features. Molecular ion species include both positive ions from proton addition $[M+H]^+$ and negative ions from proton abstraction $[M-H]^-$ as well as positive ions from alkali metal attachment and negative ions from acetate or chloride attachment. Collisional activation of both MALDI and ESI behave very similarly in that generated molecular species yield product ions that reveal many structural features of the fatty acyl lipids that can be detected in tandem mass spectrometric experiments. For many lipid species, collision induced dissociation of the positive $[M+H]^+$ reveals information about the polar headgroup, while collision induced dissociation of the negative $[M-H]^-$ provides information about the fatty acyl chain. The mechanisms of formation of many of these lipid product ions have been studied in detail and many established pathways are reviewed here. Specific examples of mass spectrometric behavior of several molecular species are presented, including fatty acids, triacylglycerol, phosphatidic acid, phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, phosphatidylserine, phosphatidylglycerol, ceramide, and sphingomyelin. © 2010 Wiley Periodicals, Inc., Mass Spec Rev 30:579–599, 2011

Keywords: triacylglycerol; phosphatidic acid; phosphatidylcholine; phosphatidylethanolamine; phosphatidylinositol; phosphatidylserine; phosphatidylglycerol; ceramide; sphingomyelin; phospholipids; neutral lipids; collisional activation; sphingolipids

I. INTRODUCTION

Lipids are a diverse array of biomolecules that are essential for all living and replicating cells. Since comprehensive discussion of the mass spectrometry of all known lipid species is not practical,

this review will focus on lipid families that are most commonly the target of mass spectrometric analysis. These species tend to have long fatty acyl chains and result from thioester carbanion condensations catalyzed by fatty acid synthase.

Analysis of these molecules by mass spectrometry can be challenging from many different points of view depending upon the type of lipid and its abundance within a biological extract. While modern mass spectrometers and the ionization methods of electrospray ionization (ESI) and matrix-assisted laser desorption ionization (MALDI) are capable of generating molecular ion species from all known lipids, the biochemistry that leads to their production often results in the generation of many closely related molecules. This is especially true for those lipids that incorporate long hydrocarbon chains derived from fatty acid biosynthesis. Successful analysis of these molecules requires considerable attention to isolation and purification in addition to consideration of mass spectrometric parameters.

Long-chain lipid substances have been recently categorized into four different families (Table 1) depending upon their mechanism of biosynthesis as well as the number of species within each family (Fahy et al., 2005). Even though this division is somewhat arbitrary, it has been useful because mass spectrometric methods developed for lipid analysis have centered around these divisions. The glycerolipids (triacylglycerols (TAGs) and diacylglycerols (DAGs)) could logically be placed in the same group as the phospholipids (PLs) (phosphatidylcholine, phosphatidylethanolamine, etc.) since both family members share a common glycerol backbone in their structure. However, these two types of lipids have entirely different chemical properties (neutral vs. polar) and biochemical functions (energy storage vs. membrane bilayer structure). In addition, living systems have vast numbers of molecular species in each of these lipid families.

The complexity alluded to above is due to the number of closely related species, often termed molecular species, which are synthesized by cells. However, not all lipids are present in cells as families of molecular species. For example, mammalian cells produce cholesterol as a single, very abundant sterol making both quantitative and qualitative analyses fairly straightforward. In contrast, the PLs, glycerolipids, and sphingolipids that are the subject of this review exist as hundreds if not thousands of different molecular species within a single mammalian cell. These molecular species differ by fatty acyl or fatty alkyl ether substituents covalently linked to a common backbone moiety such as a long-chain base (sphinganine) or glycerol as well as the positions of double bonds and acylation position of each fatty acid. This chemical diversity greatly complicates both their qualitative as well as quantitative analysis.

Another challenging aspect to lipid analysis is that some exceedingly important lipid species are only present in very low

Contract grant sponsor: National Institute of General Medical Sciences, National Institutes of Health; Contract grant number: GM069338; Contract grant sponsor: National Institute on Aging, National Institute of Health; Contract grant number: AG20238.

*Correspondence to: Robert C. Murphy, Department of Pharmacology, University of Colorado Denver, Mail Stop 8303, RC1 South, L18-6120, 12801 E. 17th Avenue, Aurora, CO 80045.

E-mail: robert.murphy@ucdenver.edu

**Correspondence to: Paul H. Axelsen, Department of Pharmacology, Biochemistry and Biophysics, University of Pennsylvania School of Medicine, Room 105 Johnson Pavilion, 3610 Hamilton Walk, Philadelphia, PA 19104-6084. E-mail: axe@upenn.edu

TABLE 1. Classification of lipids containing long hydrocarbon chains derived from the biosynthesis of fatty acids and examples of common roles played by these lipid species

		Cellular Function
Fatty Acids		
Fatty acids		Energy production, intermediates
Fatty acids amides		Signaling
Prostaglandins		Signaling
Glycerolipids		
Triacylglycerols		Energy storage
Diacylglycerols		Signaling, intermediates
Monoetherdiacylglycerols		Intermediates
Phospholipids		
Phosphatidic acid		Intermediates, signaling
Glycerolphosphoethanolamine lipids		Membrane bilayer
Glycerolphosphocholine lipids		Membrane bilayer
Glycerolphosphoserine lipids		Membrane bilayer
Glycerolphosphoinositol lipids		Membrane bilayer
Glycerolphosphoglycerol lipids		Membrane bilayer
Cardiolipin		Mitochondria bilayer
Sphingolipids		
Sphingomyelin		Membrane bilayer, microdomains
Ceramides		Signaling
Gangliosides		Protein modulation
Sulfatides		Adhesive molecule

concentrations. For example, several lipid species serve either as signaling molecules that are recognized by specific proteins, or as intermediates in enzymatic transformations. These lipids are typically present at picomolar or lower concentrations within cells or biological fluids. Another example is lipids that have been the target of reactive oxygen attack. These oxidized lipids often have important functions but are minor in abundance relative to their precursor pool. When these specific lipids are the target of mass spectrometric analysis, they present a considerable challenge because of their trace concentrations, the presence of isomeric forms, and the chromatographic separation techniques that are required.

Many types of mass spectrometric instrumentation have been employed with success in the analysis of lipids. ESI is useful for continuous analysis of liquid chromatographic effluents. It readily generates positive ions, although negative ion abundance is higher for many different lipids. Electron ionization (EI) is typically used in combination with on-line gas chromatography and prior hydrolysis and derivatization. It is a generally applicable ionization method that yields positive ions with rich fragmentation patterns (Murphy, 1993). Chemical ionization (CI) has also been used in lipid analysis in tandem with gas chromatographic separation. Atmospheric pressure chemical ionization (APCI) is similar to ESI in concept, except that charge is applied to the analyte by ion-molecule reactions during evaporation of the carrier solvent. This is a more energetic form of ionization that frequently induces fragmentation “in-source” through collisions with gas at atmospheric pressure. The fragmentation of ions in-source can be desirable and useful for

lipid analysis, as described below. Fast atom bombardment (FAB) has been used effectively in the past (Griffiths, 2003) but has largely been supplanted by ESI, MALDI, and APCI.

The most widely used mass analyzer for lipid studies has been the tandem quadrupole “MS/MS” or MS^2 mass spectrometer. This design is popular because it facilitates fragmentation by collisional activation or “collision-induced decomposition” (CID) of molecular ions in conjunction with precursor ion scanning, product ion scanning, neutral loss scanning, and multiple reaction monitoring (MRM). Each of these operating modes has valuable applications in lipid analysis. Ion traps, whether an ion cyclotron resonance instrument, octupole, or linear quadrupole design, have also been used with success for lipid analysis but only for molecular ion detection or product ion analysis. This same limitation also applies to time-of-flight (TOF) instruments that are often coupled to MALDI sources. More complex TOF instruments may be configured with an ESI source and quadrupoles for ion selection and CID (QqTOF). These instruments offer enhanced sensitivity and mass determinations that are sufficiently accurate, for example, to distinguish between lipids containing a vinyl ether group ($-\text{CH}=\text{CH}-\text{O}-$, 42.0111 Da) and an alkyl chain ($-\text{CH}_2-\text{CH}_2-\text{CH}_2$, 42.0481 Da). However, they are not as versatile as the tandem quadrupole mass spectrometer for the scanning and MRM analyses mentioned above.

Throughout this review, M will be used to designate the mass of an uncharged molecule, with the loss of a proton to create an anion denoted by $[M - \text{H}]^-$ and the gain of a proton to create a cation denoted by $[M + \text{H}]^+$. Integral masses or element symbols will be included within the brackets to indicate various other losses and gains (e.g., $[M + \text{H} - 18]^+$ to indicate the loss of water or $[M + \text{Li}]^+$ to indicate the gain of a lithium ion). The quantity actually measured by a mass spectrometer will be denoted by the mass-to-charge ratio, or m/z . Fatty acyl chains, individually and collectively within a lipid molecule, may be described with TC:DB notation in which TC is the total number of carbon atoms and DB is the total number of double bonds in the chains.

II. LIPID EXTRACTION AND IONIZATION

Textbooks generally suggest that lipids are soluble in organic solvents. This statement is true for some but not for all lipids. There are very polar lipids that have little or no solubility in organic solvents and therefore cannot be extracted out of an aqueous environment by solvent systems such as chloroform/methanol. Nonetheless, solvent extractions such as the Folch extraction (Folch, Lees, & Stanley, 1957) and the Bligh/Dyer extraction (Bligh & Dyer, 1959) are quite effective in isolation of most PLs, neutral lipids, and sphingolipids. Newer extraction solvents systems continue to be reported (Matyash, Liebisch, & Kurzchalia, 2008) and one should never underestimate the importance of this first extraction step to isolate relatively pure lipids for subsequent analysis.

There is also value in selective solvent extraction strategies to simplify lipid mixtures prior to chromatographic separation. Triglycerides and cholesterol esters can be selectively extracted with organic solvents such as isooctane/ethyl acetate (Hutchins, Barkley, & Murphy, 2008). Careful consideration and development of a specific extraction strategy can profoundly affect the rapidity of analysis and the robustness of the overall protocol. For

example, an advantageous aspect of the Bligh/Dyer (Bligh & Dyer, 1959) procedure is that formation of a single organic/aqueous phase (monophase) is an intermediate step in this extraction protocol, prior to phase separation. At this monophase step, equilibrium mixing of internal standards with the endogenous lipid substances is greatly facilitated. Some lipids are best recovered in the monophase rather than separating into two phases, for example, certain sphingolipids rather equally distribute in the upper and lower phase.

Other methods to isolate target lipids include solid-phase extraction approaches that capitalize on the hydrophobic character of the target lipid to associate with the covalently bonded hydrophobic moiety, such as an 18-carbon alkyl chain on a silica bead. An advantage of solid phase extraction is that a polar functionality, which may prevent organic solvent extraction by reducing the partition of the entire lipid into the immiscible organic phase, does not interfere with the hydrophobic interaction of the non-polar region of the lipid structure with the solid-phase material. This interaction permits facile separation of lipophilic material from non-lipophilic material.

While not the subject of this tutorial, separation techniques play a critical role in lipid analysis. Both normal phase and reversed phase liquid chromatography can be directly coupled to mass spectrometry through an ESI source (commonly denoted as "LC/MS" or "ESI-LC/MS"). Much of the classical research in lipid biochemistry involved normal phase separation of lipids using thin-layer chromatography (Touchstone, 1995), and the fundamental principles of this separation technique may be implemented with the coupling of normal phase HPLC (NP-HPLC) to the mass spectrometer. In general, normal phase chromatography separates lipids according to their polar features, whereas reversed phase HPLC (RP-HPLC) separates according to their hydrophobic properties. Either way, an important aspect of separation by HPLC prior to mass spectrometry is the removal of materials that have the potential for ion suppression during ESI.

III. MASS SPECTRAL CHARACTERISTICS

A. Fatty Acids and Eicosanoids

Fatty acids (FA) are a basic building block of many complex lipids and are found in biological extracts not only as free carboxylic acids but also esterified as in wax esters, PLs, DAGs, and TAGs (Fig. 1). They are also found as simple amides and as complex amides N-linked to the long-chain bases of sphingolipids. There is great structural diversity in this class including unsaturated, branched, hydroxyl, keto, and di-carboxylic acids. A rich history exists for the use of mass spectrometry applied to the analysis of free carboxylic acids and their volatile derivatives using EI (Murphy, 1993). As an example, the EI mass spectrum of methyl arachidate (methyl-20:0) (Fig. 1A) and methyl arachidonate (methyl-20:4) (Fig. 1B) reveal both the power and weakness of this ionization technique. For saturated fatty acids, a great deal of structural information is generated by EI including an abundant and characteristic rearrangement ion (m/z 74, the McLafferty rearrangement ion), a molecular ion at m/z 326, and various fragment ions. The ion chemistry of saturated fatty acid derivatives has been extensively studied for decades and summarized in monographs (Murphy, 1993; Pittenauer & Welz, 2006). Polyunsaturated fatty acids, in contrast, are extensively

fragmented by EI, suggesting multiple sites of radical cation formation. Typically, very little specific information is obtained from the EI mass spectrum including little or no molecular ion abundance.

Fatty acid analysis by EI is typically carried out by GC/MS and where GC retention time contributes important information to assign structural features such as methyl group branching site, *cis/trans* double bond geometry, double bond location, and other structural features by comparison with known fatty acids and their gas chromatographic elution order. Only a little information about these structural features can be obtained by interpreting the mass spectrum itself. Specific ion chemistry after CI with reactive reagent ions (Lawrence & Brenna, 2006) as well as specialized derivatization techniques, which lead to well behaved rearrangement ions during the ionization process (Harvey, 1998), have also been successful in revealing double bond location.

One approach to the analysis of complex lipids is to chemically degrade them, releasing the acyl-ester linked, *N*-acyl linked, or O-alkyl-ether linked residues that confer species identity for collective analysis. The goal of this analysis may be simple quantitation but may also include establishing the number and position of double bonds, the number and position of oxygen-containing functional groups, and alkyl branch points. However, it is important to note that a lot of information about the chemical nature of a complex lipid is lost when it is degraded in this way.

As a class, alkyl carboxylic acids have pK_a values in water between 4 and 5. In aqueous solution they tend to form micelles or bind to membranes or proteins in biological systems, and these circumstances will alter their pK_a . Nevertheless, in most cases they are anions at neutral or physiological pH. Positive mode ESI is greatly facilitated by the presence of metal ions and the formation of cationic "metalated" adducts. These adducts can be made to decompose by collisional activation in the mass spectrometer with alkyl chain C-C bond cleavage at high collision energies. The position of the double bond in monounsaturated species, for example, may be revealed by a relatively small number of fragmentation products involving cleavage at the double bond or either of the two adjacent single bonds. A better strategy has been to collisionally activate $[M + \text{metal}]^+$ ions in a high-energy instrument, which can induce site remote fragmentation and reveal double bond positions (Cheng & Gross, 2000).

Fatty acids are more typically analyzed by ESI as negative ions. As with most all $[M - H]^-$ anions of carboxylic acids, collisional activation causes free fatty acids to decompose with a neutral loss of water to yield an $[M - H - 18]^-$ ion. Somewhat less frequently, decomposition occurs via neutral loss of CO_2 to yield an $[M - H - 44]^-$ ion. Abundant carboxylate anions are formed in ESI due to the solution stability of this ionic form and these ions can be used to unambiguously establish molecular weight of the fatty acid (Fig. 1C,D). While RP-HPLC retention times are useful in establishing certain structural features, this chromatographic technique is somewhat less powerful when compared to GC, due to the reduced number of theoretical plates that HPLC separation can achieve. Collisional activation of $[M - H]^-$ ions generated as fatty acid carboxylate anions does yield some product ions from polyunsaturated FA (Fig. 1D), but little decomposition is observed from saturated FA (Fig. 1C). The combined tools of GC/MS with EI and tandem mass spectrometry of ESI generated $[M - H]^-$ ions, and $[M + \text{metal}]^+$ ions present a powerful approach for structural characterization of FA even when present in complex mixtures.




FIGURE 1. EI mass spectra of fatty acid methyl esters and collision-induced decomposition (CID) of carboxylate anions from free carboxylic acids. **A:** EI mass spectrum of methyl arachidate (70 eV); **(B)** EI mass spectrum of methyl arachidonate (70 eV); **(C)** CID mass spectrum of the carboxylate anion derived from arachidonic acid by ESI in the negative ion mode; **(D)** CID mass spectrum of the carboxylate anion derived from arachidonic acid by ESI in the negative ion mode. The CID mass spectra for this figure and subsequent figures were obtained using a tandem quadrupole (ABI 4000 Qtrap).

Fatty acid (FA) quantitation has been successfully performed with a variety of approaches, including EI-GC/MS, ESI-LC/MS, and ESI-LC/MS/MS. For such studies, stable isotope dilution is the preferred method for quantitative assessment of a target FA (Watson & Sparkman, 2007). Alternatively, a homologous internal standard may be used for those instances that do not require a high level of quantitative certainty or when isotope-labeled internal standards are not available. Quantitation is based on an empirically determined standard curve that converts the signal derived from a target molecule into an absolute quantity. This involves performing a linear regression analysis of the analyte/standard ion yield ratio versus the analyte/standard molar concentration ratio (Fig. 2). This example presents the quantitation of a metabolite of arachidonic acid generated by 5-lipoxygenase (Murphy & Gijon, 2007) and is typical of an eicosanoid standard curve used to measure such lipid mediators.

Perhaps the widest application of FA analysis by mass spectrometry has been in the study of biologically active arachidonic acid metabolites. These 20-carbon “eicosanoid”

metabolites include the prostaglandins, leukotrienes, and other complex lipids derived from cyclooxygenase, lipoxygenase, cytochrome P₄₅₀, as well as those derived from other enzymatic reactions that generate bioactive molecules from this unique polyunsaturated fatty acid (Smith & Murphy, 2008). In addition to the enzymatically formed eicosanoids, there are also products formed by free radical oxidation of polyunsaturated fatty acids esterified to PLs. When arachidonate is the target of a non-enzymatic free radical reaction, a family of compounds called “isoprostanes” are formed. These compounds are isomers of the prostaglandins formed enzymatically, and some have prostaglandin-like biological activity. Isoprostanes have been widely used as biomarkers of oxidative stress (Milne, Yin, & Morrow, 2008).

To use mass spectrometry for the quantitative analysis of eicosanoids, some unique parameter must also be included in the analytical protocol because isobaric molecules exist in many cases and molecular weight information alone is not sufficient to distinguish among them. This additional parameter could involve chromatographic elution at a specific retention time, or unique




FIGURE 2. Quantitation by stable isotope dilution of lipids typically involves establishing a standard curve where a fixed amount of a deuterium-labeled internal standard (in this case d_5 -LTC₄) is added to increasing quantities of unlabeled LTC₄. When the area for the MRM transition for LTC₄ (m/z 624 → 272) is divided by the area for the corresponding internal standard (m/z 629 → 272 for d_5 -LTC₄), a linear relationship is found with an excellent correlation coefficient.

ion chemistry following collisional activation (a specific mass loss from the molecular ion) or often both parameters. To achieve requisite sensitivity for GC/MS analysis, negative ion CI is employed after derivatization to the pentafluorobenzyl ester (PFB). Negative ion CI of this electronegative derivative yields a highly abundant $[M - \text{PFB}]^-$ carboxylate anion (Blair, 1990). Gas chromatography of these esters can separate many of the isomeric isoprostanes and all of the primary prostaglandins from each other. Many deuterium-labeled internal standards are commercially available for prostaglandin and leukotriene analysis.

A second approach is to use LC/MS/MS technology where carboxylate $[M - \text{H}]^-$ anions are directly formed by ESI, then collisionally activated to yield specific product ions. These product ions and the precursor molecular ion species can be used as a highly specific MRM experiment that is well suited for quantitative analysis. Since modern instrumentation permits the MRM experiment to be performed on numerous selected ion transitions, often as many as 50 different MRMs in a single analytical protocol, the analysis of eicosanoids in a biological sample may be carried out for a large number of both cyclooxygenase and lipoxygenase metabolites in a single run, greatly reducing the cost per analyte. These assays typically have limits of detection in picograms injected on column (Masoodi et al., 2008). A recent review has summarized our current understanding of the mechanistic origin of product ions from most of the known metabolites of arachidonate acid following collisional activation. Such knowledge is essential in developing either a qualitative or quantitative assay for these types of fatty acids (Murphy et al., 2005).

B. Glycerolipids (GL)

The analysis of glycerolipids, including TAGs and DAGs that have no electrostatic charge in solution, can nevertheless be carried out with APCI, ESI, and in many cases, EI-GC/MS


(Fig. 3). Most glycerol lipids yield a molecular ion of very low abundance when EI is employed. Abundant fragment ions corresponding to loss of each fatty acyl group as carboxyl radical and abundant low mass ions specific for each fatty acyl chain are observed (Murphy, 1993). Thus, it is difficult to determine their mass directly using EI alone, especially when mixtures of TAGs are present. When APCI conditions are employed, fairly weak $[M + \text{H}]^+$ ions are observed, but ions corresponding to the CID loss of fatty acyl groups esterified to the glycerol backbone as a neutral carboxylic acid $[M + \text{H} - \text{R}_{1,2,3}\text{COOH}]^+$ are abundant (Byrdwell, 2005). Since APCI is a relatively energetic ionization technique, the molecular ion species formed in the atmospheric pressure region of the ion source undergo CID prior to entering the low pressure of the mass spectrometer.

The analysis of TAG molecular species analysis can be challenging if information about all unique molecular species is desired. This is because of the sheer number of these species arising from the seemingly random assembly of the fatty acyl groups on the glycerol framework. For example, more than 10 different TAG species contain 52 carbon atoms and 4 double bonds in their fatty acyl chains (i.e., 52:4), and are all observed at the same mass, namely m/z 854.7364. In addition, there are positional isomers and variations in double bond geometry or location (McAnoy, Wu, & Murphy, 2005). ESI generates abundant $[M + \text{NH}_4]^+$ or $[M + \text{metal}]^+$ for TAG species when ammonium ions or alkali metal ions are present in the solvent system being sprayed (Hsu & Turk, 1999; Han & Gross, 2001).

To determine the nature of the three groups in each TAG molecular species, a second phase of CID (MS/MS/MS or MS^3) may be employed (McAnoy, Wu, & Murphy, 2005). MS/MS of $[M + \text{NH}_4]^+$ ions generates three “diglyceride type ions” for each isobaric TAG molecular species (for reactions in this section, R₁, R₂, and R₃ correspond to long-chain groups at the sn-1, sn-2, and sn-3 positions of glycerol)



These ions correspond to the loss of one esterified fatty acyl group as the free carboxylic acid as well as ammonia (reactions 1a–1c) from the molecular ion, and they provide information about the fatty acyl groups of the TAG molecular species. A mechanism responsible for the formation of these ions has been suggested (Scheme 1) (McAnoy, Wu, & Murphy, 2005). Each of these diglyceride ions can then be further decomposed in a second stage of CID to create “monoglyceride” ions. The observed mass-to-charge ratio of this ion reveals the nature of a second fatty acyl group. The difference in mass (neutral loss) between the “diglyceride ion” and the “monoglyceride ion” reveals the third fatty acyl group (reactions 2a–2d).






FIGURE 3. **A:** ESI in the positive ion mode of synthetic 17:1/14:0/18:1 triacylglycerol in the presence of electrospray buffer containing 5 mM ammonium acetate; **(B)** CID of $[M + \text{NH}_4]^+$ at m/z 834 resulting in indicated product ions corresponding to loss of each of the fatty acyl substituents as a free carboxylate anion.


The structure of the $[R_3\text{CH}_2\text{CO}]^+$ and $[R_3\text{CH}_2\text{CO} + 74]^+$ ions are indicated in Scheme 2. Chromatographic separation of TAG molecular species has been described for simple mixtures from biological sources that have relatively few common fatty acids in their lipidome (Dugo et al., 2006). For mammalian cells, complete separation of all TAG species has not been possible with either RP-HPLC or even the more powerful capillary GC columns. One is often forced to compromise and quantify the isobaric molecular species (Hutchins, Barkley, & Murphy, 2008).

At the level of molecular ion analysis, it is possible to eliminate the systematic error due to naturally abundant carbon-13 through simple calculations (Han & Gross, 2001). Nevertheless, important assumptions are made because of the influence of double bonds and ion stability of each of the different molecular species, and their influence on absolute quantitation relative to molecular ion abundance (Li & Evans, 2005). However, if the data were used in comparative experiments where the same cell population or tissue sample is differently treated, then these errors largely cancel out. Even without absolute concentrations, relative changes in the signals arising from isobaric ion species (after isotope correction) can reveal a


wealth of information about changes in this class of lipids as a result of pharmacological intervention. When an interesting change in the abundance of an ion is observed, MS/MS and MS³ may be employed to drill down and provide more specific information about the molecular species that comprise a set of isobaric ions.

One note of caution regarding assigning TAG isobaric species identity: while it is possible to calculate TC:DB for an observed mass-to-charge ratio (several computer programs do this), built into this calculation is an assumption that the observed mass-to-charge ratio is derived from a TAG. However, it is now known that in mammalian systems, monoalkyldiacylglycerols exist, which have the same nominal mass as TAGs containing one odd chain fatty acyl group. Fortunately, these unique triacylglycerol molecular species can be separated by normal phase HPLC because of the difference in polarity between an ether group and an ester group, and separation removes the assumption concerning the nature of the triacylglycerol (Hutchins, Barkley, & Murphy, 2008).

When ESI is employed, abundant and specific ions can be generated for DAGs if the free hydroxyl group is derivatized.



SCHEME 1



SCHEME 2

Derivatization may be implemented easily on the microchemistry scale and can yield excellent results for DAG analysis. Several reports have appeared using this strategy (Callender et al., 2007; Li et al., 2007).

A similar derivatization strategy has been used to analyze DAGs by GC/MS where thermostability and volatility of the lipid is an absolute requirement. This approach can be readily implemented and used to effect chromatographic separation of diverse molecular species (Hubbard et al., 1996).

C. Phospholipids (PL)

The bilayer membranes in and around all living cells are composed of large part of glycerophospholipids (PL) (Fig. 4). The generic structure of these molecules is that of a glycerol backbone where the carbinol oxygen atoms corresponding to carbon-1 and carbon-2 (commonly named according to the stereospecific numbering convention as the *sn*-1 and *sn*-2 positions) are linked to hydrophobic groups, typically by esterification to long-chain fatty acids (Fig. 4), although in some cases via alkyl ether linkages to position *sn*-1. The third (*sn*-3) carbinol is linked to a phosphate group either as a monoester (phosphatidic acid, PA) or as a phosphodiester which in turn is linked to a polar alcohol including choline (phosphatidylcho-

lines, PC), ethanolamine (phosphatidylethanolamines, PE), serine (phosphatidylserines, PS), glycerol (phosphatidylglycerols, PG), and myoinositol (phosphatidylinositols, PI). PA, PC, PE, PS, PG, and PI comprise the six major PL headgroup classes. Within a cell the diversity of this family of lipids can include a thousand or more molecular species that differ in hydrophobic groups linked to the *sn*-1 and *sn*-2 positions as well as their polar head groups. This diversity in molecular species arises from *de novo* PL biosynthesis via the Kennedy pathway (Kennedy & Weiss, 1956; Weiss, Smith, & Kennedy, 1958) as well as the active remodeling of fatty acyl groups via the Lands pathway (Lands, 1960).

Both ESI and MALDI generate $[M + H]^+$ and $[M - H]^-$ ions for PLs, although the polar head group generally drives a preference for either positive ion (PC) or negative ion formation (PS, PI, PA, and PG). PE yields both abundant positive and negative molecular ion species by either ionization method. The abundance of these molecular ion species is often used to indicate the relative quantity of each component in a mixture.

There are important differences between the nominal mass of a molecular ion (calculated with integer mass for the most abundant stable isotopes), its exact mass (calculated with the exact mass on the physical atomic scale of the most abundant stable isotopes), and its chemical molecular weight. For example, 16:0/18:1-PC has a nominal mass of 759 Da, whereas its exact mass is 759.578 Da, and its chemical molecular weight is 760.076 g/mol. The term “mass defect” refers to the difference between the nominal mass and the exact mass, and it is due to the non-integral masses of elements other than carbon. The number




FIGURE 4. General structures of glycerophospholipids (**left**) and sphingolipids (**right**). For the glycerophospholipids, stereospecific sites on the glycerol backbone are numbered (“sn” numbering). R₁ and R₂ represent alkyl chains that may be saturated or unsaturated. Together with an attached —CH₂CO— group, they constitute the “fatty acyl” chains at positions *sn*-1 and *sn*-2 of the glycerol backbone. The α methylene groups of these chains are depicted apart from the R groups because their hydrogen atoms participate in some of the decomposition reactions discussed in the text. The —CH₂CO— group at *sn*-1 position is replaced with —O— in ether lipids, and with —CH=CH—O— in vinyl ether lipids or plasmalogens. The R₃ group defines the six major phospholipid headgroup classes. For sphingolipids, R₄ is usually —C₁₃H₂₇ but may be —C₁₂H₂₅. The R₅ is an alkyl group ranging in length from 2 to 20 carbon atoms. R₆ may be —H or —OH.


of hydrogen atoms in a molecule is largely responsible for the mass defect because the exact mass of a hydrogen atom is 1.00783 Da. Chemical molecular weight differs from exact mass due to the presence of naturally abundant isotopes, chiefly carbon-13. The natural abundance of the carbon-13 isotope changes the quantity of total ionization carried by the mono-isotopic $[M + H]^+$ ion (i.e., the ion in which all carbon atoms are carbon-12). This leads to a natural trend in which the fractional abundance of a molecular species appearing as the mono-isotopic ion decreases as molecular weight increases. For example, 16:0/18:1-PC, which has 42 carbons, yields a mono-isotopic ion at *m/z* 760.586 that represents 61.7% of the ions created when counting all stable isotopes. The remaining 38.3% appears in carbon-13 and oxygen-18 containing ions at *M* + 1, *M* + 2, etc. This mono-isotopic abundance mole fraction decreases to 48.7% for 18:2/18:2/18:2/18:2-cardiolipin observed at *m/z* 1447.965. Correction for this isotope abundance variation is often made to correct the mole fraction of molecular species in a complex mixture of PL isolated from biological systems when the data correspond to the abundance of the mono-isotopic molecular ion species (Han & Gross, 2001).

Another complication in the analysis of PLs is that $[M + H]^+$ or $[M - H]^-$ ions may represent isobaric molecular species with different fatty acyl groups esterified to the glycerol carbinols. The identity of the fatty acyl groups can be determined by collisional activation of the $[M - H]^-$ ions to generate diagnostic fatty acyl ions (see below). For example, lipid anions at a single *m/z* commonly fragment to yield two carboxylate anions at relatively high abundance and additional carboxylate anions of lower abundance derived from isomeric PLs also present. One way to circumvent confusion with this issue is to report the number of carbons and the number of double bonds in the fatty acyl chains in the TC:DB notation, without stipulating which fatty acyl groups are actually present.




1. MS Identification and Analysis of PLs

Qualitative and quantitative analyses of specific PLs require some insight into their patterns of CID. A general pattern is that collisional activation of $[M + H]^+$ ions tends to yield information about their polar head groups, whereas collisional activation of $[M - H]^-$ ions reveals information about specific fatty acyl or alkylether chains present on the glycerol backbone. PLs can also form adducts with a variety of metal ions to yield both cations and anions. In many cases, these adducts increase the information content of positive ion experiments concerning acyl/alkylether chains, and of negative ion experiments concerning polar headgroups (Han & Gross, 2005).

Protonated PL cations undergo relatively few informative decompositions under normal circumstances. Aside from the headgroup-specific ions described below, the only ion consistently observed in positive mode experiments results from the concerted neutral losses of the phospho-headgroup and the *sn*-2 chain as a ketene (reaction 3a; Scheme 3; Fig. 5A). De-protonated PL anions, in contrast, undergo several common and informative CID for all six PL headgroup classes. The simplest decompositions are charged and neutral losses of the phospho-headgroup (reactions 3b and 3c). The relative yield of these reactions is chiefly determined by the ability of the headgroup to stabilize a negative charge.



Another common negative ion decomposition produces fatty acid anions from the *sn*-1 and *sn*-2 chains (reactions 4a and





SCHEME 3

4b; Scheme 3). The mechanism of reactions 4a and 4b appears deceptively simple; however, the relative yields from these two reactions vary widely among the various PL classes. These differences in yield constitute evidence that the CID behavior involves some degree of interaction between headgroups and acyl chains. At minimum, this interaction involves migration of charge from the phosphate group, although other interactions are likely to be involved and will be discussed below in separate discussions of each PL class. The relative yields of reactions 4a and 4b may also vary with collision energy, possibly due to the effect of this parameter on rates of secondary ion decomposition rather than on rates of ion production.





Either of the acyl chains may also leave as a neutral fatty acid with the retention of charge on the lyso-PL remnant (reactions 5a and 5b; Scheme 4). These reactions provide an indirect means to ascertain the chemical composition of the *sn*-1 and *sn*-2 chains. The remnants may undergo additional concerted decomposition reactions involving the headgroup and release the other chain as a fatty acid anion (reactions 5c and 5d; Scheme 4). Analysis of such remnants is also the most direct way to determine the composition of ether-linked *sn*-1 substituents, because these chains cannot be released as carboxylate anions.




Finally, the neutral loss of acyl chains may occur via ketene formation (reactions 6a and 6b). Compared to the neutral loss of a fatty acid, the neutral loss of a ketene leaves an –OH group on the charged lyso-PL remnant (Scheme 5).

Oftentimes, collisional activation results in a single decomposition reaction, and the product ion does not retain sufficient internal energy for additional decompositions. The –OH group remaining after the loss of a ketene, however, may facilitate concerted decomposition reactions in an MS^2 experiment that produce a phospho-headgroup ion (reaction 6c), or charged cyclic phosphoglycerol remnants (reactions 6d and 6e; Scheme 6). The loss of both chains as ketenes leaves –OH groups on both the *sn*-1 and *sn*-2 positions and an ion at $[R_3 + 170]^-$ that is often observed in abundance. MS^3 experiments have shown that this ion decomposes into an $[R_3 + 96]^-$ ion (reaction 6f; Scheme 7), but since it may also arise via reaction 3b or 6c (Scheme 6), its origin in an MS^2 experiment is not certain.




2. Patterns Characteristic of PA Lipids

Phosphatidic acid (PA) lipids have the simplest structures among PL classes, but exchangeable protons on the terminal phosphate group in these lipids facilitate several complex reactions that are uncommon in other PL classes (Hsu & Turk, 2000c) (Fig. 5). In




SCHEME 4

positive mode analyses, the most important decomposition is heterolytic cleavage of a polar C–O bond with neutral loss of H_3PO_4 and the formation of an $[M - 97]^{+}$ glyceride cation that is diagnostic of PA lipids (reaction 7, Scheme 8; Fig. 5A). An $[R_1 + 116]^{+}$ ion is also formed according to reaction 3a.




The intrinsic negative charge of PA lipids facilitates detection of the parent $[M - H]^-$ ion in negative mode analyses,



SCHEME 5


usually making it much more abundant than the $[M + H]^{+}$ ion in positive mode analyses. In addition, the array of ions produced by CID is more diverse and informative in negative mode analyses. For example, cleavage of a polar O–P bond produces charged loss of the phosphate group as PO_3^- (m/z 79) (reaction 8). Less commonly, the C–O bond is cleaved and the phosphate is lost as $H_2PO_4^-$ (m/z 97) (reaction 9)



Either fatty acyl chain may leave as a charged species via reactions 4a and 4b, as neutral acids via reactions 5a and 5b, or as ketenes via reactions 6a and 6b (Fig. 5B). The loss of one chain as a ketene provides a proximate hydroxyl group that facilitates the concerted loss of the other chain as an acid, yielding a major ion with m/z 153 (reactions 6d and 6e; Scheme 6). The neutral loss of one chain as a fatty acid according to reactions 4a and 4b yields a major $[M - R_1CH_2COOH - H]^-$ ion. In-source CID studies have demonstrated that this ion loses 136 Da to produce a fatty acid anion (reactions 5c and 5d; Scheme 4 with $R_3 = H$). However, the extent to which these reactions add to the yield of reactions 4a and 4b in an MS/MS experiment is unclear.

3. Patterns Characteristic of PE Lipids


Condensation of ethanolamine and glycerophosphate yields a zwitterionic lipid headgroup with distinctive CID patterns (Fig. 6). In positive mode analyses, the $[M + H]^{+}$ ion might be expected to form by protonation of the phosphate, since it bears a formal negative charge, and protonation of the basic amino group to form a cationic site. Fragmentation at the C–O bond by mechanisms analogous to those described above for PA lipids would be expected to then yield a phosphoethanolamine cation appearing at m/z 142 (Scheme 9). However, this scenario is not




SCHEME 6

observed (Fig. 6A). Instead, there is neutral loss of phosphoethanolamine, and the proton associates with an ester group to form $[M - 140]^+$ ions that are diagnostic of lipids in the PE class (reaction 10), but of lower abundance for ether lipids and plasmalogens (Zemski-Berry & Murphy, 2004). As with PA lipids, an $[R_1 + 116]^+$ ion may be observed. A richer array of


positive ions yielding information about chain identities may be obtained from lithium ion adducts of PE (Hsu & Turk, 2000a).



Negative mode CID of PE lipids (Fig. 6B) produces weak phosphoethanolamine anions at m/z 140 according to reaction 3b, and abundant fatty acid anions according to reactions 4a and 4b (Hsu & Turk, 2000e). Neutral chain losses occur according to reactions 5a and 5b and 6a and 6b, with the most abundant



SCHEME 7



SCHEME 8




FIGURE 5. Positive and negative ion CID mass spectra from 1-heptadecanoyl-2-arachidonoyl-3-glycerophosphate (17:0/20:4-PA). Molecular ion species generated by ESI. **A:** Collisional activation of the $[M + H]^+$ ion at m/z 711.5; **(B)** collisional activation of the $[M - H]^-$ ion at m/z 709.5.

observed ion corresponding to loss of a ketene from the *sn*-2 fatty acyl group in the example shown (Fig. 6B).

4. Patterns Characteristic of PC Lipids

Lipids with the PC headgroup exhibit distinctive CID behavior due to the permanent positive charge of the quaternary nitrogen (Fig. 7). All species yield abundant molecular ion species in positive ion mode studies, most likely because the phosphate group readily accepts a proton during ESI. Molecular ion species decompose efficiently to produce phosphocholine ions with m/z 184 that are diagnostic for the PC headgroup class (reaction 11, Fig. 7). However, these ions do not appear to be produced by simple cleavage of the C–O bond at *sn*-3 (Haroldsen & Gaskell, 1989). Instead, stable isotope studies indicate that a proton is acquired from the β carbon of the *sn*-2 chain (Scheme 10). Because phosphocholine ions are produced with high efficiency, other decomposition products are only produced in relatively low abundance (Fig. 7A).




FIGURE 6. Positive and negative ion CID mass spectra from 1-heptadecanoyl-2-arachidonoyl-3-glycerophosphoethanolamine (17:0/20:4-PE). Molecular ion species generated by ESI. **A:** Collisional activation of the $[M + H]^+$ ion at m/z 754.5; **(B)** collisional activation of the $[M - H]^-$ ion at m/z 752.5.

The phosphate group in PC lipids also readily accepts alkali metal ions, and a distribution of species containing Na^+ and K^+ can complicate quantitative analysis in biological samples. Metal ions also induce additional decomposition patterns. For example, sodiated PC lipids produce $[M - 59 + \text{Na}]^+$ ions representing the neutral loss of $(\text{CH}_3)_3\text{N}$, and $[M - 183 + \text{Na}]^+$ ions representing the neutral loss of phosphocholine (Ho & Huang, 2002). An ability to detect the PC headgroup by the neutral loss of phosphocholine rather than the production of phosphocholine cations is advantageous when using ion trap instrumentation, because it is easier to trap ions with m/z values that are closer to the parent ion. In some cases, metal ions induce enough neutral loss of RCOOH and $\text{RCH}=\text{C=O}$ to identify the acyl chains (Ho & Huang, 2002).

Phosphatidylcholine (PC) lipids form negative ions by forming adducts with anions such as chloride $[M + 35]^-$ and acetate $[M + 59]^-$ (Fig. 7B). The anions are loosely bound and may be lost before entering the spectrometer at high orifice (de-clustering) potentials (Harrison & Murphy, 1995). On the other hand, when removed they often result in the neutral loss of a methyl group and the formation of $[M - 15]^-$ ions at the orifice.





FIGURE 7. Positive and negative ion CID mass spectra from 1-heptadecanoyl-2-arachidonoyl-3-glycerophosphocholine (17:0/20:4-PC). Molecular ion species generated by ESI. **(A)**: Collisional activation of the $[M + H]^+$ ion at m/z 796.6; **(B)**: collisional activation of the $[M + \text{acetate}]^-$ ion at m/z 854.6.


Thus, by judicious choice of solvent conditions and orifice potentials, anion adduct as well as $[M - 15]^-$ ions are available for the analysis by negative mode mass spectrometry. When subjected to collisional activation, these ions yield decomposition products according to reactions 4–6 (Hsu & Turk, 2003).

5. Patterns Characteristic of PS Lipids

Despite their negative charge at neutral pH, lipids with the PS headgroup readily form $[M + H]^+$ ions (Hsu & Turk, 2005) (Fig. 8). They decompose to $[M - 184]^+$ ions by neutral loss of phosphoserine with such efficiency that few other ions are produced (reaction 12a; Scheme 11; Fig. 8A). Nevertheless, an $[R_1 + 116]^+$ ion according to reaction 3a may be observed, and secondary fragmentations may occur to produce acylium ions from the fatty acyl chains (Scheme 11).

Phosphatidylserine (PS) lipids may also bind up to three alkali cations. $[M + \text{Li}]^+$ ions decompose into $[M - 184]^+$ ions as with protonated species; however, they also produce lithium–phosphoserine adduct ions with m/z 192 and undergo the neutral loss of serine to produce $[M - 87]^+$ ions (Hsu & Turk, 2005). Lithiated-PS may undergo rearrangement during decomposition to produce an $[M + \text{Li} - 98]^+$ ion with an apparent TAG structure. Neutral losses from these ions enable identification of the *sn*-1 and *sn*-2 chains. By virtue of their two anionic sites, PS lipids

may bind two or three Li^+ ions to form $[M - \text{H} + 2\text{Li}]^+$ and $[M - 2\text{H} + 3\text{Li}]^+$. The decomposition products obtained are analogous to those obtained from $[M + \text{Li}]^+$ ions, although there are marked differences in yields. In most cases, Na^+ may substitute for Li^+ , with altered product ion yields.





$[M - H]^-$ ions readily decompose to $[M - 88]^-$ ions from the neutral loss of serine, often occurring in-source (Fig. 8B; reaction 12b; Scheme 12). Isotope exchange studies suggest that the mechanism of this decomposition involves a hydrogen from either the amino or the carboxyl groups of the serine (Hsu & Turk, 2005). The resulting $[M - 88]^-$ ions are equivalent to the $[M - H]^-$ ions of PA, and they decompose accordingly within an MS^2 experiment to yield fatty acid anions and remnant ions following the neutral loss of fatty acid chains or ketenes (Hsu & Turk, 2005). PS lipids may form lithiated $[M - 2H + Li]^-$ ions, or their sodiated analog. These ions follow qualitatively similar decomposition patterns as $[M - H]^-$ ions, but with an increased tendency to decompose by neutral loss of ketenes.

6. Patterns Characteristic of PG Lipids

Phosphatidylglycerol (PG) lipids form $[M + H]^+$ ions that decompose via neutral loss of glycerophosphate to yield $[M - 171]^+$ product ions that are diagnostic of the PG headgroup






FIGURE 8. Positive and negative ion CID mass spectra from 1-heptadecanoyl-2-arachidonoyl-3-glycerophosphoserine (17:0/20:4-PS). Molecular ion species generated by ESI. **A:** Collisional activation of the $[M + H]^+$ ion at m/z 798.5; **(B)** Collisional activation of the $[M - H]^-$ ion at m/z 796.5.


(Fig. 9A; reaction 13; Scheme 13) (Fig. 9). They also produce an $[R_1 + 116]^+$ ion according to reaction 3a.



Collision-induced negative mode decomposition of PG $[M - H]^-$ ions yields several informative ions via reactions 4–6 (Fig. 9B) (Hsu & Turk, 2001). Secondary decompositions within an MS^2 experiment are also common, facilitated and influenced by the free hydroxyl groups of the glycerol. For example, one chain may leave as $RCOOH$, and the other may leave as a ketene, to yield an intermediate ion with m/z 227 (reactions 14a and 14b; small signal present in Fig. 9B). The glycerol residue in this ion appears to undergo the neutral loss of $C_3H_6O_2$ to yield a product with m/z 153 (reaction 6f; Scheme 14; Fig. 9B). PG lipids that have lost an acyl chain as $RCOOH$ or $RCH=C=O$ may also undergo neutral losses of $C_3H_6O_2$ via reaction 6f to yield $[M - RCOOH - 75]^-$ ($[R_1 + 191]^-$, m/z 405 in Fig. 9B; reaction 14c; Scheme 14) or $[M - RCH=C=O - 75]^-$ ions (Hsu & Turk, 2001).



SCHEME 11






SCHEME 12



7. Patterns Characteristic of PI Lipids

Phosphatidylinositol (PI) lipids can form positive ions (Fig. 10A) that decompose upon collisional activation by the loss of phosphoinositol (reaction 15a; Scheme 15) (Fig. 10). However, negative ions are more abundant and more informative due to the intrinsic negative charge of the PI headgroup (Fig. 10B). PI lipid anions decompose according to reactions 3–6 and under some conditions, inositol may be lost as a neutral fragment to yield the corresponding PA species (Hsu & Turk, 2000b).



SCHEME 13

The close proximity of hydroxyl groups on the inositol ring to the phosphate facilitates concerted decomposition reactions involving two or more CID reactions in a single MS^2 experiment. For example, reaction 3b yields m/z 259 (reaction 15b), but the concomitant loss of H_2O , presumably by the condensation of proximate hydroxyl groups on phosphate and inositol, produces an even more abundant m/z 241 ion (Scheme 16; Fig. 10B). Another neutral loss of the inositol ring in the same MS^2 experiment accounts for a major ion that is observed at m/z 153 (Hsu & Turk, 2000b). Yet another example of three concerted reactions in an MS^2 experiment begins with the loss of two neutral fatty acids to produce an m/z 297 ion. Adjacent hydroxyl groups then appear to promote cleavage of an O–P bond and lose $C_3H_5O_2$ to produce an m/z 223 ion that is prominent in MS^2 spectra (Scheme 17). PI lipids may have one (PIP) or two (PIP₂) additional phosphate groups on the inositol ring, each of which may add another negative charge to the molecular ion. Because the molecular ion species and their decomposition products may be multiply charged, the mass spectra become quite complex to interpret. Their decomposition mechanisms are similar to those of PI lipids; however, the phosphate groups may also undergo various gas-phase rearrangements after CID that produce additional ions such as diphosphate (Hsu & Turk, 2000b).

8. Patterns Characteristic of Cardiolipins

The general structure of cardiolipins shares all of its motifs with PA lipids, and therefore they undergo the same CID reactions (Hsu et al., 2005). However, their spectrum is complicated by the presence of two phosphate groups, three glycerol backbones, and four fatty acyl chains per molecule, along with the tendency to



A**B**


FIGURE 9. Positive and negative ion CID mass spectra from 1-heptadecanoyl-2-arachidonoyl-3-glycerophosphoglycerol (17:0/20:4-PG). Molecular ion species generated by ESI. **A:** Collisional activation of the $[M + H]^+$ ion at m/z 785.5; **(B)** collisional activation of the $[M - H]^-$ ion at m/z 783.5.

bear two negative charges and thereby become doubly charged. Therefore, cardiolipins may form various ions that combine two or more of the motifs observed with PA lipids. One such ion is produced following the concerted neutral loss of two fatty acid chains from one glycerol (Scheme 18). The resulting ion may then lose a 136 Da fragment to produce an ion equivalent to an $[M - H]^-$ PA ion.

D. Sphingolipids

Sphingolipids are perhaps the most diverse long-chain lipids found in cells in terms of chemical structure and biological function (Fig. 4). While these derivatives of long-chain bases such as sphingosine and sphinganine were originally thought to be primarily located at the outer membrane bilayer of cells, it is now known that they have intracellular roles as well as serving as mediators in intracellular signaling (Table 1) (Merrill et al., 2007). The diversity in their structure is also reflected in their chemical properties and approaches used by mass spectrometry to analyze these molecules.

$[M - H - RCH_2COOH]^-$
(product of reactions 5a,b)



SCHEME 14

Efficient isolation of sphingolipids and preparation for mass spectrometric analysis can be quite challenging. Some sphingolipids do not favorably distribute in organic extraction systems and can be found in equal concentrations in both the aqueous as well as the organic layers of a chloroform/methanol/water extraction protocol. In fact, one approach has been to analyze the monophase of the Bligh/Dyer extract method to maximize the yield of the sphingolipids targeted for analysis (Sullards et al., 2007). One widely used technique to isolate sphingolipids relies on their stability to high pH. Glycerolipids and glycerophospholipids are co-extracted, but their ester groups are saponified to free fatty acids and glycerol at high pH, leaving the sphingolipids concentrated in the organic extract.

The analysis of sphingolipids by mass spectrometry can be exemplified in approaches to quantify signaling ceramides (Merrill et al., 2005), analysis of sphingomyelin (a PL incorporating a long-chain base as well as an *N*-fatty acyl chain), and glycosphingolipids (Merrill et al., 2005). The diversity in mass spectrometric behavior of the latter group of sphingolipids is best studied in context of complex carbohydrate analysis since structural challenges revolve around the carbohydrate subunit identity and linkage analysis.





FIGURE 10. Positive and negative ion CID mass spectra from 1-heptadecanoyl-2-arachidonoyl-3-glycerophosphoinositol (17:0/20:4-PI). Molecular ion species generated by ESI. **A:** Collisional activation of the $[M + H]^+$ ion at m/z 873.5; **(B)** collisional activation of the $[M - H]^-$ ion at m/z 871.5.

1. Ceramides


Ceramides are the simplest long-chain sphingolipids, consisting of a fatty acyl chain N-linked to a long-chain base known as sphingosine (Figs. 4 and 11). However, their analysis is complicated by variations in carbon chain length, position and degree of unsaturation, and position and number of hydroxylation sites. The most abundant ceramides have an 18-carbon sphingosine and an acyl chain of variable length. They may or may not have a double bond in the acyl chain and a hydroxyl group on the second carbon of the acyl chain. In positive mode analysis (Fig. 11A), common ceramides readily lose H_2O to form $[\text{M} - \text{H}_2\text{O} + \text{H}]^+$ ions and undergo amide bond cleavage with neutral loss of the fatty acid to yield an ion with m/z 264 (Scheme 19) (Fillet et al., 2002). This has been a particularly useful product ion for MRM analysis of ceramides (Merrill et al., 2007).

The decomposition pattern of ceramide anions is particularly rich and varies in an informative way depending on structural variation of the long-chain base and the fatty acyl substituent such as a 2'-hydroxyl ceramide (Han, 2002; Hsu &


Turk, 2002). For example, $[M - H - 327]^-$ ions are produced only when the hydroxyl group is present in the N-fatty acyl substituent (Scheme 20). A very abundant CID product ion corresponds to $[M - H - 256]^-$ which most likely is a conjugated anion (Scheme 20). This same decomposition pathway involving an initial loss of water can lead to the abundant ion at m/z 237. Another common product ion is $[M - H - 240]^-$ which is abundant in Figure 11B.

2. *Sphingomyelin*


Sphingomyelin is a ceramide bearing an O-linked phosphocholine group (Figs. 4 and 12). The efficiency with which collisionally activated sphingomyelin produces phosphocholine ions (in a manner similar to that illustrated in Scheme 10) is sufficiently high that very few other cations are produced with ESI (Fig. 12A). Some investigators have produced an array of more informative ions by first producing ceramide cations in-source with APCI (Karlsson, Michelsen, & Odham, 1998). Others have produced a similar array of product ions by ESI of lithium adducts (Hsu & Turk, 2000d). As with sodium-PE adducts, lithiated-sphingomyelin $[M + Li]^+$ ions produce $[M - 59 + Li]^+$ ions representing the neutral loss of $(CH_3)_3N$. $[M - 183 + Li]^+$



SCHEME 16





SCHEME 18




SCHEME 17

ions representing the neutral loss of phosphocholine, and $[M - 429]^+$ representing the neutral losses of phosphocholine and most of the sphingosine chain.

Sphingomyelin produces ions for negative mode mass spectrometry by forming anion adducts in a manner similar to that of PC lipids (Fig. 12B). However, sphingomyelin has no acyl-linked chains and therefore, they do not produce informative decomposition products with simple collisional activation. This problem may be solved by selecting anion adducts that produce $[M - 15]^-$ ions by *N*-demethylation, and subjecting the product ions to further collisional activation in an MS^3 experiment. Under these conditions, neutral loss of the *N*-acyl chain is observed, permitting one to distinguish SM from PC, and identify the *N*-acyl and long-chain base substituents (Houjou et al., 2004).



SCHEME 19



SCHEME 20




FIGURE 12. Positive and negative ion CID mass spectra from *N*-dodecanoyl-sphingomyelin (SM d18:1/12:0). Molecular ion species generated by ESI. **A:** Collisional activation of the $[M + H]^+$ ion at m/z 647.5; **(B)** collisional activation of the $[M + \text{acetate}]^-$ ion at m/z 705.5.

3. Glycosphingolipids

Glycosphingolipids (GSLs) are ceramides with one or more O-linked sugar residues that may be modified by phosphate, sulfate, or various other groups. In positive mode analysis, relatively low collision energies cause neutral loss of the glycoside with retention of the charge on the ceramide. Higher energies in positive mode and negative mode afford insight into the glycoside composition. The nomenclature of GSLs and their decompensation patterns is complex and has been reviewed by others (Levery, 2005).

ACKNOWLEDGMENTS

The PL standards were gifts from Avanti Polar Lipids. This work was supported in part by a grant (LIPID MAPS) from the National Institute of General Medical Sciences of the National Institutes of Health to R.C.M. (GM069338), and from the National Institute on Aging of the National Institutes of Health to P.H.A. (AG20238). The authors also acknowledge Christopher Johnson and Charis Uhlson for the lipid mass spectra and Robert Barkley for critical reading of the manuscript.

REFERENCES

Blair IA. 1990. Electron-capture negative-ion chemical ionization mass spectrometry of lipid mediators. *Methods Enzymol* 187:13–23.

Bligh EG, Dyer WJ. 1959. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37:911–917.

Byrdwell WC. 2005. The bottom-up solution to the triacylglycerol lipidome using atmospheric pressure chemical ionization mass spectrometry. *Lipids* 340:383–417.

Callender HL, Forrester JS, Ivanova P, Preininger A, Milne S, Brown HA. 2007. Quantification of diacylglycerol species from cellular extracts by electrospray ionization mass spectrometry using a linear regression algorithm. *Anal Chem* 79:263–272.

Cheng C, Gross ML. 2000. Applications and mechanisms of charge-remote fragmentation. *Mass Spectrom Rev* 19:398–420.

Dugo P, Kumm T, Crupi ML, Cotroneo A, Mondello L. 2006. Comprehensive two-dimensional liquid chromatography combined with mass spectrometric detection in the analyses of triacylglycerols in natural lipidic matrixes. *J Chromatogr A* 1112:269–275.

Fahy E, Subramaniam S, Brown HA, Glass CK, Merrill AH, Jr., Murphy RC, Raetz CRH, Russell DW, Seyama Y, Shaw W, Shimizu T, Spener F, van Meer G, VanNieuwenhze MS, White SH, Witztum JL, Dennis EA. 2005. A comprehensive classification system for lipids. *J Lipid Res* 46:839–862.

Fillet M, Van Heugen JC, Servais AC, De Graeve J, Crommen J. 2002. Separation, identification and quantitation of ceramides in human cancer cells by liquid chromatography-electrospray ionization tandem mass spectrometry. *J Chromatogr A* 949:225–233.

Folch J, Lees M, Stanley GHS. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226:497.

Griffiths WJ. 2003. Tandem mass spectrometry in the study of fatty acids, bile acids and steroids. *Mass Spectrom Rev* 22:81–152.

Han XL. 2002. Characterization and direct quantitation of ceramide molecular species from lipid extracts of biological samples by electrospray ionization tandem mass spectrometry. *Anal Biochem* 302:199–212.

Han X, Gross RW. 2001. Quantitative analysis and molecular species fingerprinting of triacylglyceride molecular species directly from lipid extracts of biological samples by electrospray ionization tandem mass spectrometry. *Anal Biochem* 295:88–100.

Han X, Gross RW. 2005. Shotgun lipidomics: Electrospray ionization mass spectrometric analysis and quantitation of cellular lipidomes directly from crude extracts of biological samples. *Mass Spectrom Rev* 24:367–412.

Haroldsen PE, Gaskell SJ. 1989. Quantitative analysis of platelet activating factor using fast atom bombardment/tandem mass spectrometry. *Biomed Environ Mass Spectrom* 18:439–444.

Harrison KA, Murphy RC. 1995. Negative electrospray ionization of glycerophosphocholine lipids: Formation of $[M - 15]^-$ ions occurs via collisional decomposition of adduct anions. *J Mass Spectrom* 30:1772–1773.

Harvey DJ. 1998. Picolinyl esters for the structural determination of fatty acids by GC/MS. *Mol Biotechnol* 10:251–260.

Ho YP, Huang PC. 2002. A novel structural analysis of glycerophosphocholines as TFA/K⁺ adducts by electrospray ionization ion trap tandem mass spectrometry. *Rapid Commun Mass Spectrom* 16:1582–1589.

Houjou T, Yamatani K, Nakanishi H, Imagawa M, Shimizu T, Taguchi R. 2004. Rapid and selective identification of molecular species in phosphatidylcholine and sphingomyelin by conditional neutral loss scanning and MS^3 . *Rapid Commun Mass Spectrom* 18:3123–3130.

Hsu FF, Turk J. 1999. Structural characterization of triacylglycerols as lithiated adducts by electrospray ionization mass spectrometry using low-energy collisionally activated dissociation on a triple stage quadrupole instrument. *J Am Soc Mass Spectrom* 10:587–599.

Hsu FF, Turk J. 2000a. Characterization of phosphatidylethanolamine as a lithiated adduct by triple quadrupole tandem mass spectrometry with electrospray ionization. *J Mass Spectrom* 35:596–606.

Hsu FF, Turk J. 2000b. Characterization of phosphatidylinositol, phosphatidylinositol-4-phosphate, and phosphatidylinositol-4,5-bisphosphate

by electrospray ionization tandem mass spectrometry: A mechanistic study. *J Am Soc Mass Spectrom* 11:986–999.

Hsu FF, Turk J. 2000c. Charge-driven fragmentation processes in diacyl glycerophosphatidic acids upon low-energy collisional activation. A mechanistic proposal. *J Am Soc Mass Spectrom* 11:797–803.

Hsu FF, Turk J. 2000d. Structural determination of sphingomyelin by tandem mass spectrometry with electrospray ionization. *J Am Soc Mass Spectrom* 11:437–449.

Hsu FF, Turk J. 2000e. Charge-remote and charge-driven fragmentation processes in diacyl glycerophosphoethanolamine upon low-energy collisional activation: A mechanistic proposal. *J Am Soc Mass Spectrom* 11:892–899.

Hsu FF, Turk J. 2001. Studies on phosphatidylglycerol with triple quadrupole tandem mass spectrometry with electrospray ionization: Fragmentation processes and structural characterization. *J Am Soc Mass Spectrom* 12:1036–1043.

Hsu FF, Turk J. 2002. Characterization of ceramides by low energy collisional-activated dissociation tandem mass spectrometry with negative-ion electrospray ionization. *J Am Soc Mass Spectrom* 13: 558–570.

Hsu FF, Turk J. 2003. Electrospray ionization/tandem quadrupole mass spectrometric studies on phosphatidylcholines: The fragmentation processes. *J Am Soc Mass Spectrom* 14:352–363.

Hsu FF, Turk J. 2005. Studies on phosphatidylserine by tandem quadrupole and multiple stage quadrupole ion-trap mass spectrometry with electrospray ionization: Structural characterization and the fragmentation processes. *J Am Soc Mass Spectrom* 16:1510–1522.

Hsu FF, Turk J, Rhoades ER, Russell DG, Shi YX, Groisman EA. 2005. Structural characterization of cardiolipin by tandem quadrupole and multiple-stage quadrupole ion-trap mass spectrometry with electrospray ionization. *J Am Soc Mass Spectrom* 16:491–504.

Hubbard WC, Hundley TR, Oriente A, MacGlashan DW, Jr. 1996. Quantitation of 1-stearoyl-2-arachidonoyl-sn-3-glycerol in human basophils via gas chromatography-negative ion chemical ionization mass spectrometry. *Anal Biochem* 236:309–321.

Hutchins PM, Barkley RM, Murphy RC. 2008. Separation of cellular non-polar neutral lipids by normal phase chromatography and analysis by electrospray ionization mass spectrometry. *J Lipid Res* 49:804–813.

Karlsson Å, Michelsen P, Odham G. 1998. Molecular species of sphingomyelin: Determination by high-performance liquid chromatography/mass spectrometry with electrospray and high performance liquid chromatography/tandem mass spectrometry with atmospheric pressure chemical ionization. *J Mass Spectrom* 33:1192–1198.

Kennedy EP, Weiss SB. 1956. Function of cytidine coenzymes in the biosynthesis of phospholipids. *J Biol Chem* 222:193–213.

Lands WEM. 1960. Metabolism of glycerolipids 2. Enzymatic acylation of lysolecithin. *J Biol Chem* 235:2233–2237.

Lawrence P, Brenna JT. 2006. Acetonitrile covalent adduct chemical ionization mass spectrometry for double bond localization in non-methylene-interrupted polyene fatty acid methyl esters. *Anal Chem* 78:1312–1317.

Levery SB. 2005. Glycosphingolipid structural analysis and glycosphingolipidomics. *Methods Enzymol* 405:300–369.

Li X, Evans JJ. 2005. Examining the collision-induced decomposition spectra of ammoniated triglycerides as a function of fatty acid chain length and degree of unsaturation. I. The OXO/YOY series. *Rapid Commun Mass Spectrom* 19:2528–2538.

Li YL, Su X, Stahl PD, Gross ML. 2007. Quantification of diacylglycerol molecular species in biological samples by electrospray ionization mass spectrometry after one-step derivatization. *Anal Chem* 79:1569–1574.

Masoodi M, Mir AA, Petasis NA, Serhan CN, Nicolaou A. 2008. Simultaneous lipidomic analysis of three families of bioactive lipid mediators leukotrienes, resolvins, protectins and related hydroxy-fatty acids by liquid chromatography/electrospray ionisation tandem mass spectrometry. *Rapid Commun Mass Spectrom* 22:75–83.

Matyash V, Liebisch G, Kurzchalia TV. 2008. Lipid extraction by methyl-tert-butyl ether for high-throughput lipidomics. *J Lipid Res* 49:1137–1146.

McAnoy AM, Wu CC, Murphy RC. 2005. Direct qualitative analysis of triacylglycerols by electrospray mass spectrometry using a linear ion trap. *J Am Soc Mass Spectrom* 16:1498–1509.

Merrill AH, Jr., Sullards MC, Allegood JC, Kelly S, Wang E. 2005. Sphingolipidomics: High-throughput, structure-specific, and quantitative analysis of sphingolipids by liquid chromatography tandem mass spectrometry. *Methods* 36:207–224.

Merrill AH, Jr., Wang MD, Park M, Sullards MC. 2007. (Glyco)sphingolipidology: An amazing challenge and opportunity for systems biology. *Trends Biochem Sci* 32:457–468.

Milne GL, Yin H, Morrow JD. 2008. Human biochemistry of the isoprostanate pathway. *J Biol Chem* 283:15533–15537.

Murphy RC. 1993. Mass spectrometry of lipids. New York: Plenum Press.

Murphy RC, Gijon MA. 2007. Biosynthesis and metabolism of leukotrienes. *Biochem J* 405:379–395.

Murphy RC, Barkley RM, Zemski Berry K, Hankin JA, Harrison K, Johnson C, Krank J, McAnoy A, Uhlsom C, Zarini S. 2005. Electrospray ionization and tandem mass spectrometry of eicosanoids. *Anal Biochem* 346:1–42.

Pitterauer E, Welz W. 2006. Free fatty acids and triacylglycerols. In: Caprioli RM, editor. *Biological applications, Part B: Carbohydrates, nucleic acids and other biological compounds*. Oxford, UK: Elsevier. pp 397–415.

Smith WL, Murphy RC. 2008. The eicosanoids: Cyclooxygenase, lipoxygenase, and epoxygenase pathways. In: Vance DE, Vance JE, editors. *Biochemistry of lipids, lipoproteins & membranes*. Oxford, UK: Elsevier Science. pp 331–362.

Sullards MC, Allegood JC, Kelly S, Wang E, Haynes CA, Park H, Chen Y, Merrill AH, Jr. 2007. Structure-specific, quantitative methods for analysis of sphingolipids by liquid chromatography-tandem mass spectrometry: "Inside-out" sphingolipidomics. *Methods Enzymol* 432:83–115.

Touchstone JC. 1995. Thin-layer chromatographic procedures for lipid separation. *J Chromatogr B Biomed Appl* 671:169–195.

Watson JT, Sparkman OD. 2007. *Introduction to mass spectrometry: Instrumentation, applications, and strategies for data interpretation*. Chichester, England: Wiley & Sons.

Weiss SB, Smith SW, Kennedy EP. 1958. Enzymatic formation of lecithin from cytidine diphosphate choline and D-1,2-diglyceride. *J Biol Chem* 231:53–71.

Zemski-Berry KA, Murphy RC. 2004. Electrospray ionization tandem mass spectrometry of glycerophosphoethanolamine plasmalogen phospholipids. *J Am Soc Mass Spectrom* 15:1499–1508.